

2026年7月BLASTAMによるいもち病の発病条件

病害虫防除所

日付	岐阜	大垣	愛西 海津	関ヶ 原	揖斐 川	樽見 根尾	美濃	八幡	長滝 白鳥	美濃 加茂	黒川 白川	多治 見	中津 川	恵那	萩原	宮地 下呂	金山	高山	白川 村	六所 庄川	宮之 前朝日	河合	神岡	栃尾 上宝
7/1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7/2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	△	-	-	-
7/3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	●	-	-	●	◎	△	-	-	△
7/4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7/5	●	●	●	●	●	●	●	●	●	●	●	●	●	●	◎	●	●	-	-	-	-	-	-	-
7/6	●	-	●	-	-	-	-	-	●	●	●	-	●	●	●	●	-	●	●	△	△	●	●	△
7/7	●	-	-	-	-	●	-	●	●	●	-	-	-	-	●	●	●	-	-	△	△	-	-	△
7/8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7/9																								
7/10																								
7/11																								
7/12																								
7/13																								
7/14																								
7/15																								
7/16																								
7/17																								
7/18																								
7/19																								
7/20																								
7/21																								
7/22																								
7/23																								
7/24																								
7/25																								
7/26																								
7/27																								
7/28																								
7/29																								
7/30																								
7/31																								

- 凡例 △ : 準好適条件 (湿潤時間10時間以上だが、前5日間の平均気温は20℃未満)
 ▲ : 準好適条件 (湿潤時間10時間以上だが、前5日間の平均気温は25℃以上)
 ○ : 準好適条件 (湿潤時間10時間以上だが、湿潤期間の平均気温は15~25℃の範囲外)
 ◎ : 準好適条件 (湿潤期間が湿潤時間中の平均気温ごとの必要な時間より短い)
 ● : 好適条件 (湿潤時間中の平均気温が15℃~25℃ であり、湿潤時間が湿潤時間中の平均気温ごとに必要な時間を満たし、当日を含めてその日以前5日間の日平均気温の平均値が20℃~25℃の範囲にある)
 - : 好適条件、準好適条件は満たされなかった
 空白 : 欠測値があるため判定不能
 *感染好適条件が現れると感染し、潜伏期間 (5~10日) を経て発病します。